arXiv:1304.2040v1 [quant-ph] 7 Apr 2013

Comment on "Separable states with unique decompositions"

R. Augusiak,[1] J. Bae,[1,2] J. Tura,[1] and M. Lewenstein[1,3]
[1]*ICFO–Institut de Ciències Fotòniques, 08860 Castelldefels (Barcelona), Spain*
[2]*Center for Quantum Technologies, National University of Singapore, 3 Science Drive 2, Singapore 117543*
[3]*ICREA–Institució Catalana de Recerca i Estudis Avançats, Lluis Companys 23, 08010 Barcelona, Spain*

In the recent paper [K.-C. Ha and S.-H. Kye, *Separable states with unique decompositions*, arXiv:1210.1088], Ha and Kye claim to find a decomposable optimal entanglement witness that disproves the conjecture that structural physical approximations to optimal entanglement witnesses are separable states. It is the main aim of this note to show, both analytically and numerically, that the presented witness is not optimal, and as such is not a counterexample to the above conjecture.

Few years ago, some of us posed the following conjecture [1]: *Structural physical aproximations to optimal positive maps are entanglement breaking channels.* Via the Choi-Jamiołkowski [2] isomorphism it can be rephrased as follows. Let W be a normalized entanglement witness (EW) acting on a product Hilbert space $\mathcal{H}_A\otimes\mathcal{H}_B$ of finite dimension $d_A\times d_B$, and let us consider its convex combination with the maximally mixed state, i.e., $W(p)=pW+(1-p)\mathbb{1}/d_Ad_B$ $(0\leq p\leq 1)$. Clearly, for some range of the mixing parameter p, $W(p)$ is positive, and thus a quantum state. For the largest $p=p_*$ for which this is the case, we call $W(p_*)$ the structural physical approximation (SPA) of W [3]. Then, in terms of entanglement witnesses the conjecture states that if W is optimal, then its SPA is a separable state, and was later supported by numerous examples of witnesses (see, e.g., Ref. [5, 6]), both decomposable and indecomposable, and even studied in the continuous variables case [6].

Quite surprisingly, very recently the conjecture has been disproved by Ha and Kye [4], who have provided an example of an indencomposable optimal entanglement witness, whose structural physical approximation is entangled. Still it remains unresolved in the decomposable case. In an attempt to elucidate this the same authors proposed a witness whose SPA is entangled and argue that it is optimal [7].

The main aim of this note is to show that this decomposable witness is not optimal, and therefore it is not a correct candidate for disproving the conjecture.

Before presenting our arguments, we briefly recall the notion of optimality and some basic tools allowing to check it. For this purpose, let us denote by

$$D_W=\{\rho\,|\,\mathrm{Tr}(\rho W)<0\},\tag{1}$$

and

$$P_W=\{|e,f\rangle\,|\,\langle e,f|W|e,f\rangle=0\},\tag{2}$$

the set of all entangled states detected by W, and the set of product vectors realizing zero mean value on W.

Consider now two entanglement witnesses W_1 and W_2. Following Ref. [8] we call W_1 *finer* than W_2 if $D_{W_1}\supseteq D_{W_2}$. We then say that an EW W is *optimal* if there is no other witness that is finer than W. It was proven in Ref. [8] that a given EW W is optimal iff for any $\lambda>0$ and a positive operator P acting on $\mathcal{H}_A\otimes\mathcal{H}_B$ and having support orthogonal to P_W, $W-\lambda P$ is no longer a witness, i.e., the inequality

$$\langle e,f|W-\lambda P|e,f\rangle\geq 0\tag{3}$$

does not hold for all product vectors $|e,f\rangle$. This immediately implies a sufficient condition for optimality: an EW W is optimal if P_W spans $\mathcal{H}_A\otimes\mathcal{H}_B$.

We can now pass to the decomposable EW introduced by Ha and Kye [7]. It acts on $\mathcal{H}_A\otimes\mathcal{H}_B=\mathbb{C}^3\otimes\mathbb{C}^3$ and reads

$$\begin{aligned}W_{\theta,b}&=|w_0\rangle\!\langle w_0|^{\Gamma}+\frac{1}{b}\sum_{i=1}^{3}|w_i\rangle\!\langle w_i|^{\Gamma}\\&=\begin{pmatrix}1&0&0&0&\mathrm{e}^{\mathrm{i}\theta}&0&0&0&\mathrm{e}^{-\mathrm{i}\theta}\\0&b&0&1&0&0&0&0&0\\0&0&\frac{1}{b}&0&0&0&1&0&0\\0&1&0&\frac{1}{b}&0&0&0&0&0\\\mathrm{e}^{-\mathrm{i}\theta}&0&0&0&1&0&0&0&\mathrm{e}^{\mathrm{i}\theta}\\0&0&0&0&0&b&0&1&0\\0&0&1&0&0&0&b&0&0\\0&0&0&0&0&1&0&\frac{1}{b}&0\\\mathrm{e}^{\mathrm{i}\theta}&0&0&0&\mathrm{e}^{-\mathrm{i}\theta}&0&0&0&1\end{pmatrix},\end{aligned}\tag{4}$$

with $b>0$, $b\neq 1$, $\theta\in(-\pi/3,\pi/3)$, $\theta\neq 0$. Then, Γ stands for the partial transposition with respect to the first subsystem, and the four vectors $|w_i\rangle$ are given by

$$|w_0\rangle=|00\rangle+|11\rangle+|22\rangle,\tag{5}$$

$$|w_i\rangle=b|i-1,i\rangle+\mathrm{e}^{\mathrm{i}\theta}|i,i-1\rangle\quad(i=1,2,3),\tag{6}$$

where the summation is modulo three. It straightforwardly follows from (4) that this witness has block-diagonal form $W_{\theta,b}=\Lambda_\theta+(1/b)(P_1+P_2+P_3)$, where Λ_θ is given by

$$\Lambda=\begin{pmatrix}1&\mathrm{e}^{\mathrm{i}\theta}&\mathrm{e}^{-\mathrm{i}\theta}\\\mathrm{e}^{-\mathrm{i}\theta}&1&\mathrm{e}^{\mathrm{i}\theta}\\\mathrm{e}^{\mathrm{i}\theta}&\mathrm{e}^{-\mathrm{i}\theta}&1\end{pmatrix}\tag{7}$$

and acts on on the subspace spanned by $|ii\rangle$ $(i=0,1,2)$. The three remaining blocks are rank-one matrices $P_i=|\phi_i\rangle\!\langle\phi_i|$ with $|\phi_i\rangle=b|i-1,i\rangle+|i,i-1\rangle$ $(i=1,2,3)$ being vectors spanning a subspace orthogonal to P_W, with the latter, as it is indirectly stated in Ref. [7], consisting of the following six vectors

$$\begin{aligned} |z_1^{(\pm)}\rangle &= (\pm 1, \omega^*, 0)\otimes(\omega, \mp 1, 0),\\ |z_2^{(\pm)}\rangle &= (0, \pm 1, \omega^*)\otimes(0, \omega, \mp 1),\\ |z_3^{(\pm)}\rangle &= (\omega^*, 0, \pm 1)\otimes(\mp 1, 0, \omega), \end{aligned} \tag{8}$$

where $\omega = \sqrt{b}\,\mathrm{e}^{\mathrm{i}\theta/2}$.

For any $\theta \in (-\pi/3, \pi/3)$ except for $\theta = 0$, the witness has exactly one negative eigenvalue which comes from Λ_θ, while for $\theta = 0$, Λ_0 is an unnormalized projector onto $|00\rangle + |11\rangle + |22\rangle$, and therefore $W_{0,b}$ becomes a positive matrix for any positive b.

By referring to the results of Ref. [9], but without presenting an explicit proof, Ha and Kye argue that $W_{\theta,b}$ has the spanning property. This, via the result recalled above would directly imply that it is optimal. It is later shown that for $\theta \in (-\pi/12, \pi/12)$ except for $\theta = 0$, and b satisfying some simple condition (see Ref. [7]), the SPA of $W_{\theta,b}$ is entangled. Provided that both facts are valid, this would serve as a counterexample to the conjecture.

Our aim is to comment on the first part of their reasoning and show that $W_{\theta,b}$ is not optimal for any choice of the parameters θ and b. First of all, it should be stressed that P_W consists of only six vectors (8), and therefore, contrary to what is stated in Ref. [7], $W_{\theta,b}$ cannot have the spanning property. This does not imply, nevertheless, that the witness is not optimal. It is known that there exist optimal decomposable witnesses without the spanning property [10].

In what follows we will prove that indeed the witness is not optimal, and the method we will utilize in order to reach this goal was developed in the appendix of Ref. [5]. It can be briefly summarized as follows. Let W be some decomposable witness and consider a positive number ϵ and a positive operator P, whose support is orthogonal to P_W (in particular, it can be a projector onto one of the vectors $|\phi_i\rangle$ introduced above). Clearly, for all product vectors $|e, f\rangle$ satisfying

$$\langle e, f|W|e, f\rangle > \epsilon \tag{9}$$

there always exists $\lambda > 0$ (obeying $\lambda < \epsilon/M$ with $M = \max_{|e,f\rangle}\langle e, f|P|e, f\rangle$) such that the inequality (3) is fulfilled. It is obvious that the same holds for any, even arbitrarily small but positive ϵ and the corresponding product vectors obeying (9). One then needs to check whether (3) remains satisfied by product vectors that do not obey (9), i.e., those for which $\langle e, f|W|e, f\rangle \leq \epsilon$, where, in some sense, ϵ is a free parameter that can be set arbitrarily small. In our case P_W a finite set, and a product vector $|e, f\rangle$ obeying the latter inequality must be "close" to one of elements of P_W. In other words, for any such vector $|e, f\rangle$ there is a normalized $|e_0, f_0\rangle \in P_W$ such that its local components can be written as

$$|e\rangle = \frac{1}{\sqrt{1 + |\delta|^2 + |\delta'|^2}}(|e_0\rangle + \delta|e_1\rangle + \delta'|e_2\rangle), \tag{10}$$

and

$$|f\rangle = \frac{1}{\sqrt{1 + |\varepsilon|^2 + |\varepsilon'|^2}}(|f_0\rangle + \varepsilon|f_1\rangle + \varepsilon'|f_2\rangle), \tag{11}$$

where $|e_i\rangle$ and $|f_i\rangle$ $(i = 1, 2)$ are normalized vectors orthogonal to $|e_0\rangle$ and $|f_0\rangle$, respectively, while δ, δ', ε, and ε' are complex numbers of vanishing absolute values for $\epsilon \to 0$.

With the aid of Eqs. (10) and (11), we can now write

$$\langle e, f|W|e, f\rangle = \sum_{i=0}^{4} A_i(W), \tag{12}$$

where by $A_i(W)$ we have denoted terms of the ith degree in variables δ, δ', ϵ, and ϵ'. From the fact that $|e_0, f_0\rangle \in P_W$ it follows that $A_0(W) = A_1(W) = 0$, and the first nonvanishing term in Eq. (12) can be $A_2(W)$. On the other hand, $A_3(W)$ and $A_4(W)$ can be made arbitrarily small by appropriately adjusting ϵ. It is then clear that if $A_2(W)$ is always positive except for $\delta = \delta' = \epsilon = \epsilon' = 0$ for any element of P_W, then W is optimal [8].

We can now apply the above reasoning to the witness $W_{\theta,b}$. To this end, consider a particular pair of vectors from the corresponding P_W, say $|z_1^{(\pm)}\rangle$. Up to normalization, their local components read $|e_0^{(\pm)}\rangle \propto (\pm 1, \omega^*, 0)$ and $|f_0^{(\pm)}\rangle \propto (\omega, \mp 1, 0)$, while vectors orthogonal to both of them can be taken as $|e_1^{(\pm)}\rangle \propto (\mp\omega, 1, 0)$ and $|e_2^{(\pm)}\rangle = (0, 0, 1)$, and $|f_1^{(\pm)}\rangle \propto (\pm 1, \omega, 0)$ and $|f_2^{(\pm)}\rangle = (0, 0, 1)$. Then, after some algebra one has that

$$A_2^{(\pm)}(W) = \frac{1}{N_e N_f}\left[2(|\delta|^2 + |\varepsilon|^2) + \left(b + \frac{1}{b} - 1\right)(|\delta'|^2 + |\varepsilon'|^2) \pm \frac{4\sqrt{b}}{1+b}\sin(3\theta/2)\mathrm{Im}(\delta'\varepsilon')\right], \tag{13}$$

where $N_e = 1 + |\delta|^2 + |\delta'|^2$ and $N_f = 1 + |\varepsilon|^2 + |\varepsilon'|^2$. It is easily provable that $A_2^{(\pm)}(W)$ are strictly positive unless $\delta = \delta' = \epsilon = \epsilon' = 0$. With the aid of the facts that $\mathrm{Im}z \leq |z|$ holds for any $z \in \mathbb{C}$ and $|\sin(3\theta/2)| \leq 1$, we can lower bound them in the following way

$$A_2^{(\pm)}(W) \geq \frac{1}{N_e N_f}\left[2(|\delta|^2 + |\varepsilon|^2) + \left(b + \frac{1}{b} - 1\right)(|\delta'|^2 + |\varepsilon'|^2) - \frac{4\sqrt{b}}{1+b}|\delta'\varepsilon'|\right], \tag{14}$$

which can later be rewritten as

$$A_2^{(\pm)}(W) \geq \frac{1}{N_e N_f}\left[2(|\delta|^2+|\varepsilon|^2)+\left(b+\frac{1}{b}-1\right)(|\delta'|-|\varepsilon'|)^2+2\left(b+\frac{1}{b}-1-\frac{2\sqrt{b}}{1+b}\right)|\delta'\varepsilon'|\right], \tag{15}$$

It is not difficult to convince oneself that $b+1/b-1>1$ and

$$b+\frac{1}{b}-1-\frac{2\sqrt{b}}{1+b}>0 \tag{16}$$

for any positive $b\neq 1$, and therefore the expression in square brackets in Eq. (15) is always positive except for the case when $|\delta|=|\delta'|=|\varepsilon|=|\varepsilon'|=0$, completing the proof.

Analogous reasoning can be applied to the remaining two pairs of vectors in P_W [cf. Eq. (8)]. For each of them, $|z_2^{(\pm)}\rangle$ or $|z_3^{(\pm)}\rangle$, one obtains exactly the same $A_2^{(\pm)}(W)$ as in Eq. (13). As a consequence, the witness $W_{\theta,b}$ is not optimal for any positive $b\neq 1$, and $\theta\in(-\pi/3,\pi/3)$ and $\theta\neq 0$.

With the aid of the procedure described in Ref. [11] we have also numerically studied optimality of the witness. Since the optimization over product vectors may lead to different local minima, several initial conditions, uniformly and randomly chosen have been considered, and this procedure was repeated at least 10^3 times for each choice of θ and b. As it follows from Fig. 1, the obtained results fully support the above proof. As a positive operator to be subtracted from the witness we took $\widetilde{P}_1 = P_1/(1+b^2)$ (see above for the definition of P_1). Then, as a function of $\theta\in[-\pi/3,\pi/3]$ and $b\in(0,3)$ we determined the maximal value of λ such that $\widetilde{W}(\lambda) = (\widetilde{W}_{\theta,b}-\lambda\widetilde{P}_1)/(1-\lambda)$ is an entanglement witness (see Fig. 1), where $\widetilde{W}_{\theta,b}$ is a normalized version of $W_{\theta,b}$. It should be noticed that for $\theta=0$, $W_{0,b}$ is a positive matrix and therefore one can remove the whole block $(1/b)P_1$ from it. This corresponds to the fact that for any fixed b, the function on Fig. 1 achieves its maximum for $\theta=0$.

Acknowledgments. This work is supported by the ERC Grant QUAGATUA, EU projects AQUTE and SIQS, and Spanish TOQATA. R. A. acknowledges support from the Spanish MINCIN through the Juan de la Cierva program. J. B. acknowledges National Research Foundation and Ministry of Education of Singapore.



FIG. 1. As a function of $\theta\in[-\pi/3,\pi/3]$ and $b\in(0,3)$, it is presented the numerically determined maximal value of λ such that $(\widetilde{W}_{\theta,b}-\lambda\widetilde{P}_1)/(1-\lambda)$ is a witness. Here $\widetilde{W}_{\theta,b}$ and $\widetilde{P}_1$ denote the normalized witness $W_{\theta,b}$ and the rank-one operator P_1. The visible maximum for $\theta=0$ is a consequence of the fact that $W_{0,b}$ is a positive matrix for any allowed b, and hence one can subtract the whole block $(1/b)P_1$ from it.

[1] J. K. Korbicz, M. L. Almeida, J. Bae, M. Lewenstein, and A. Acín, Phys. Rev. A **78**, 062105 (2008).
[2] A. Jamiołkowski, Rep. Math. Phys. **3**, 275 (1972); M.-D. Choi, Linear Algebra Appl. **10**, 285 (1975).
[3] P. Horodecki, Phys. Rev. A **68**, 052101 (2003).
[4] K.-C. Ha and S.-H. Kye, J. Math. Phys. **53**, 102204 (2012).
[5] D. Chruściński and J. Pytel, Phys. Rev. A **82**, 052310 (2010); D. Chruściński, J. Pytel, G. Sarbicki, *ibid.* **80**, 062314 (2009).
[6] R. Augusiak, J. Bae, Ł. Czekaj, M. Lewenstein, J. Phys. A **44**, 185308 (2011).
[7] K.-C. Ha and S.-H. Kye, *Separable states with unique decompositions*, arXiv:1210.1088.
[8] M. Lewenstein, B. Kraus, J. I. Cirac, and P. Horodecki, Phys. Rev. A **62**, 052310 (2000).
[9] S.-H. Kye, *Facial structures for various notions of positivity and applications to the theory of entanglement, preprint*, arXiv:1202.4255.
[10] R. Augusiak, G. Sarbicki, and M. Lewenstein, Phys. Rev. A **84**, 052323 (2011).
[11] J. M. Leinaas, J. Myrheim, and P. Ø. Sollid, Phys. Rev. A **81**, 062329 (2010).